



Global polarization of hyperons in Au+Au collisions at $\sqrt{s_{NN}} = 27$ GeV in the STAR experiment

Egor Alpatov (for STAR Collaboration)


National Research Nuclear University MEPhI

Outline:


- Introduction
- Global hyperon polarization
- Motivation
- The STAR experiment
- Hyperon polarization measurements
- Results
- Conclusions

Introduction

- The Quark-Gluon Plasma (QGP) formed in non-central nuclear-nuclear collisions is associated with large orbital angular momentum, that leads to vorticity in the medium
- Spin-orbit coupling aligns spin directions of produced particles with the direction of vorticity
 - -Z.-T. Liang and X.-N. Wang, PRL94, 102301 (2005)
 - S. A. Voloshin, arXiv:nucl-th/0410089
- Another possible source of particle polarization is magnetic field, created in non-central collisions in the initial stage
 - -D. Kharzeev, L. McLerran, and H. Warringa, Nucl.Phys.A803, 227 (2008)
 - -McLerran and Skokov, Nucl. Phys. A929, 184 (2014)



Vorticity



- In non-central HIC the initial collective longitudinal flow velocity depends on x:



$$\omega_y = \frac{1}{2} (\nabla \times v)_y \approx -\frac{1}{2} \frac{dv_z}{dx}$$

- For small polarization:

Becattini, Karpenko, Lisa, Upsal, Voloshin PRC95.054902 (2017)

$$P_\Lambda \approx \frac{1}{2} \frac{\omega}{T} + \frac{\mu_\Lambda B}{T}$$

$$P_{\bar{\Lambda}} \approx \frac{1}{2} \frac{\omega}{T} - \frac{\mu_\Lambda B}{T}$$



Jiang et al, PRC94 044910 (2016)

How to measure global polarization?

- Hyperons are “self-analyzing” due to weak decay properties:
 - Daughter baryons are preferentially emitted in parent spin direction


- Daughter baryons of hyperons with polarization (\vec{P}) follows the distribution:

$$\frac{dN}{d\Omega^*} = \frac{1}{4\pi} (1 + \alpha_H |\vec{P}| \cdot \widehat{\vec{p}_b^*}) = \frac{1}{4\pi} (1 + \alpha_H P \cos \theta^*)$$

- α_H - decay parameter, unique for each hyperon species
- $\widehat{\vec{p}_b^*}$ is the daughter baryon momentum in the parent frame
- Projection to the transverse plane can be measured:

$$P_H = \frac{8}{\pi \alpha_H} \frac{\langle \sin(\psi_1 - \phi_p^*) \rangle}{Res(\psi_1)}$$


- ψ_1 is the reaction plane angle
- ψ_1 and it's resolution $Res(\psi_1)$ can be calculated with spectator's signal.




Motivation

- Global polarization of Λ hyperons was measured for $\sqrt{s_{NN}} = 7.7$ -200 GeV at STAR
- P_H decreases with increasing collision energy
- Difference between P_Λ and $P_{\bar{\Lambda}}$ maybe due to B-field effect
- Theoretical calculations can quantitatively explain the energy dependence of the Λ polarization, but many of them fail to explain differential measurements
- Nowadays there is a growing interest to measure the global polarization of Λ and $\bar{\Lambda}$ produced from the decays of other particles such as Ξ .
- Ξ polarization may provide new input for global polarization and vorticity studies

STAR PRC 98, 014910 (2018)




The STAR experiment



Detectors, used in this work:

- **Time Projection Chamber**
 η : $[-1, 1]$
- **Event Plane Detector**
 η : $[2.1, 5.1]$
- **Beam-Beam Counters**
 η : $[3.3, 5.0]$



Event plane measurement

- Event plane was measured using BBCs and EPDs

$$\Psi_1 = \tan^{-1} \left(\frac{\sum w_i \sin(\phi_i)}{\sum w_i \cos(\phi_i)} \right)$$

where w_i is detector's tile ADC

- $$Res(\Psi_{1,EPD\ East}) = Res(\Psi_{1,EPD\ West}) = \frac{1}{\sqrt{\langle \cos(\Psi_{1,EPD\ East} - \Psi_{1,EPD\ West}) \rangle}}$$

- Resolution for east and west detectors combination estimated by two sub-event method
A. M. Poskanzer, S. A. Voloshin, PRC58.1671(1998)


- $Res_{EPD} > 1.5 Res_{BBC}$
- BES 1 results used BBCs event plane

BBC




- 18 (x2) tiles but only 16(x2) photomultiplier tubes
- $|\eta_{BBC}| = 3.3 - 5.0$

EPD






- 372 (x2) tiles and 372(x2) Silicon photomultipliers
- $|\eta_{EPD}| = 2.1 - 5.1$



Hyperon reconstruction

- Hyperon daughters are identified using dE/dx in TPC
- Used topology of decays to reconstruct Λ
- Ξ were reconstructed via $\Xi \rightarrow \Lambda + \pi$



Signal extraction

- The data was fitted with the following equation:



$$\langle \sin(\Delta\varphi) \rangle^{obs} = (1 - f^{Bg}(M_{inv})) \langle \sin(\Delta\varphi) \rangle^{Sg} + f^{Bg}(M_{inv}) \langle \sin(\Delta\varphi) \rangle^{Bg}$$

$$\Delta\varphi = \Psi_1 - \varphi_p^*$$


$f^{Bg}(M_{inv})$ is background fraction

- $f^{Bg}(M_{inv})$ is taken from the invariant mass distribution fit
- Assumption: background signal = 0
- Signal is scaled after extraction:

$$P_H = \frac{8}{\pi\alpha_H} \frac{\langle \sin(\psi_1 - \varphi_p^*) \rangle}{Res(\psi_1)}$$




Results



- Ξ polarization transfers to it's daughter Λ with transfer factor $C_{\Xi\Lambda}$
- Ξ polarization was measured directly and via Λ daughter
- Directly measured P_{Ξ} is comparable with $P_{\Xi \rightarrow \Lambda}$
- Weak centrality dependence of Ξ polarization within uncertainties

T. Niida, RHIC&AGS Annual Users Meeting 2020



- P_{Ξ} is consistent with P_{Λ} trend

Results from previous publications were scaled to be consistent with updated decay parameters: Zyla, P.A. and others, Review of Particle Physics

Old $\alpha_{\Lambda} = 0.642$, $\frac{\alpha_{\Lambda old}}{\alpha_{\Lambda new}} = 0.869$

Conclusions

- We presented first results of $\Xi + \bar{\Xi}$ global polarization measurements in Au+Au collisions at 27 GeV
- Direct polarization measurements are consistent with measurements via Λ daughters
- Ξ polarization is comparable to Λ polarization at 27 GeV within uncertainties
- We are looking forward to continuing this measurements at other energies

Thank you for your attention!